

The Benefits of Exercise in Breast Cancer

Milena dos Santos Barros Campos,^{1,2} Roberta Helena Fernandes Feitosa,³ Carolina Christianini Mizzaci,⁴ Maria do Rosário Toscano von Flach,⁵ Betty Janny Maia Siqueira,⁶ Luiz Eduardo Mastrocola⁴

Clínica e Hospital São Lucas, RedeD'Or São Luiz,¹ Aracaju, SE – Brazil

Divisão de Cardiologia do Hospital Universitário de Sergipe,² Aracaju, SE – Brazil

Gerência dos Sistemas de Informações Epidemiológicas – Secretaria Municipal de Saúde,³ Goiânia, GO – Brazil

Instituto Dante Pazzanese de Cardiologia,⁴ São Paulo, SP – Brazil

Fundação Bahiana para Desenvolvimento das Ciências (FBDC),⁵ Salvador, BA – Brazil

Universidade de Pernambuco,⁶ Recife, PE – Brazil

Abstract

Breast cancer is the most prevalent cancer among women, accounting for nearly 30% of all cancers, while in men, it represents only 1% of cases. Breast cancer is the main cause of death for cancer, and its incidence and mortality vary according to patients' ethnicity, geographic region, and socioeconomic status. Due to the low prevalence of breast cancer among men and the scarcity of studies in the literature, exercises have been prescribed based on extrapolations from studies on female patients. Scientific evidence has suggested beneficial effects of physical exercises on breast cancer prevention, treatment, and post-treatment. In addition to combatting sedentary behavior, it is essential to maintain a healthy body weight, limit alcohol consumption, and follow a balanced diet, rich in fruit, vegetables, grains and fibers, and limited in red meat. The effects of exercises are not restricted to breast cancer, but extend to controlling modifiable risk factors, and reducing the incidence of cardiovascular diseases, and all-cause and cardiovascular mortality.

Introduction

Breast cancer is the most incident cancer among women in the world, with approximately 2.3 million new cases in 2020, corresponding to 24.5% of all types of cancer.¹ In Brazil, apart from non-melanoma skin cancer, breast cancer is the most common among women, with the highest rates in the south and in the southeast regions.² The highest occurrence in these regions may be attributed to the highest human development index and life expectancy, high prevalence of white race, life style, later pregnancy and less children.^{3,4} For 2022, it is estimated 66,280 new cases of breast cancer in Brazil.²

Keywords

Breast Neoplasms; Exercise Movement Techniques; Exercise; Diet; Healthy

Mailing Address: Milena dos Santos Barros Campos •

Hospital São Lucas, Rede D'Or – Av. Cel. Stanley da Silveira. Postal Code 49015-400, Aracaju, SE – Brazil

E-mail: millybarros@yahoo.com

Manuscript received February 03, 2022, revised manuscript August 08, 2022, accepted September 28, 2022

DOI: <https://doi.org/10.36660/abc.20220086>

With respect to mortality, breast cancer is the main cause of death for cancer among the female population in Brazil, except for the north region, where colon cancer ranks the first. The highest mortality rates are observed in the south and southeast regions and the rates increase from the age of 40 years of age.⁵ In places where mortality rates are low, like the north of Brazil, the possibility of underdiagnosis of breast cancer cannot be excluded.

Genetic causes, including BRCA1 and BRCA2 mutations, are responsible for 5-10% of all cases of breast and ovarian cancer, with a greater contribution of environmental factors and lifestyle in the pathogenesis of these tumors.⁶ The BRCA 1 and BRCA2 genes produce tumor suppressive proteins. These proteins repair damaged DNA and thereby play a role in maintaining genetic material stability in each cell. When mutation or alteration occurs in one of these genes, the activity of the protein product can be altered, and the DNA damage may be not properly repaired. As a result, the cells are more likely to develop genetic changes that may lead to cancer development.⁷

The adoption of a healthy lifestyle is important for prevention of breast cancer, including an adequate diet (higher consumption of fruit, vegetables and whole grains, and lower consumption of red meat), weight control, reduction of alcohol intake, and regular physical exercises (PE).⁸ The effects are not restricted to the prevention, but also to disease control, since experimental studies have demonstrated influence on tumor formation kinetics, growth and recurrence.⁹

The third consensus of the World Cancer Research Fund¹⁰ and the Brazilian Society of Oncology guidelines on physical activity¹¹ address the importance of women being physically active, involving several types of physical activities, from household chores (e.g. gardening), occupational, and recreational activities, to those systematically categorized as PE, whose frequency, intensity, time and type (aerobic, resistance and combined) are determined by prescription.

Mechanisms of action of physical exercises in tumor progression

PE promote different organic and biological mechanisms that can be involved in the control of the development of several tumors. These responses originate from metabolic and hormonal changes, in addition to modulation of systemic inflammation.¹² However, the potential of directly affecting tumor progression has been recently related to

changes in vascularization and blood flow in tumors,¹³ the use of substrates by neoplastic cells, protein interactions between cancer and muscle tissue, and to the regulation of immune function by PE.¹⁴⁻¹⁶

Tumor microenvironment acts in the cooption and deviation of the action of immunoinflammatory and stromal cells.¹⁷ While the acute and transient action of lymphocytes and macrophages is a controlling and repair factor of tissue damage, chronic inflammation and macrophage infiltration into the tissue promote tumor progression.¹⁸

The development of anticancer therapy has been based on hallmarks (biological capabilities acquired by human cells during the development of tumors) proposed by Weinberg in 2000.¹⁹ Also based on these concepts, possible mechanisms of action and adaptation by which PE can influence these marks of tumor development have been studied (Figure 1).

Results of preclinical studies indicate that these molecular effects resulting from each exercise session overcome the control of hormonal factors and insulin.²⁰ During exercise, these factors act immediately on tumor metabolism, and long-term training leads to metabolic and immunogenic adaptations that contribute to slow tumor progression.²¹

Evidence that PE inhibits malignant tumor progression emerged from animal models.^{12,22,23} Recently, clinical studies have identified cellular and molecular actions that are similar to exercise in patients with cancer, including breast cancer.^{24,25} However, there are still no studies that clearly determine the clinical relevance of these findings.

Physical exercise in breast cancer prevention

One of the first and largest prospective studies on PE and breast cancer was the “Nurses Health Study”. This

study evaluated 121 701 nurses aged between 30 and 55 years, during a follow-up period of 16 years, and showed that women who engaged in moderate or vigorous physical activity for seven or more hours per week had a nearly 20% lower risk of breast cancer than those who engaged in such physical activity for less than one hour per week (relative risk 0.82; 95% confidence interval [CI] 0.70-0.97). This benefit was observed in both premenopausal and postmenopausal women.²⁶

Literature has suggested that women who exercise regularly have 10-25% lower risk of breast cancer compared with those who do not.^{27,28} This association seems to be stronger for activity sustained over the lifetime and after menopause, for women who are normal weight, have no family history of breast cancer, and are parous.²⁹ Available evidences of the impact of PE on breast cancer reduction in BRCA1 and BRCA2 mutation carriers are limited, and larger studies are warranted.³⁰

A meta-analysis published in 2013 of 31 prospective studies found a significant association between physical activity and reduction of breast cancer risk, with a combined relative risk (RR) with 95 % CI of 0.88 (0.85-0.91), highlighting the importance of prevention. Dose-response analysis suggested that the risk of breast cancer decreased by 2 % for every 25 metabolic equivalent (MET)-h/week increment in non-occupational physical activity (approximately 10 hours per week in household activities), 3 % for every 10 MET-h/week increment in recreational activity (equivalent to 4 h/week of walking in 3Km/h), and 5 % for every 2 h/week increment in moderate and vigorous recreational activity.³¹ Other studies have corroborated these results, suggesting greater risk reduction of breast cancer with higher levels of PE.^{28,32} The exact dose and the type of exercise needed to reduce breast cancer risk have not been clearly determined.




Figure 1 – Scheme of the mechanisms by which physical education can affect the hallmarks of cancer in breast cancer. Adapted from “Hallmarks of cancer”, by BioRender.com

In an umbrella review³³ about physical activity and cancer incidence and mortality, the results suggested a reduction of breast cancer risk in the general population. However, classification of physical activity across studies were heterogeneous and most reviews were based on observational studies, mainly cohort studies, in which the control of selection bias is difficult, since healthy habits tend to cluster. For example, a person with a healthy lifestyle eat well, has close to ideal body weight and is nonsmoker. To minimize limitations of this type of study, guidelines have been established based on cohort studies with larger samples of the population.³³

The role of PE in breast cancer prevention seems to be linked to reductions in estrogen activity, insulin resistance, inflammation, and oxidative stress.³⁴ Estrogen is related to reduction of cellular proliferation and tumor development. PE increase sex hormone-binding globulin, and reduce circulating levels of estrogen, as described in Figure 1. They contribute to the reduction of fat mass, mainly by visceral fat reduction, improvement of cellular insulin sensitivity and consequent decrease of insulin serum levels. Insulin is involved in the activation of aromatase and estrogen elevation, in addition to exerting mitogenic effects. In addition, PE have immunomodulatory effects, increasing both innate and acquired immunity and improving DNA repair mechanisms, thereby reducing the risk of breast cancer.³⁵

More research is needed to fully understand the mechanisms by which physical activity can reduce breast cancer risk.

General recommendations for physical activity in the prevention and control of breast cancer

In 2020, the World Health Organization (WHO) and the Brazilian Ministry of Health recommended for the general adult population (18-64 years old) and for breast cancer

survivors, 150-300 minutes of moderate-intensity physical activity, at least 75-150 minutes of vigorous-intensity aerobic physical activity; or an equivalent combination of moderate- and vigorous-intensity activity throughout the week.^{11,36,37}

Table 1 summarizes the description of different intensities of physical activity based on the Physical Activity Guidelines for the Brazilian Population.

It is important to differ physical activities (voluntary body movements, with energy expenditure resting levels) from PE (planned, structured, and repetitive physical activity, essentially aimed to improve cardiorespiratory fitness, strength, flexibility and balance). It is recommended that PE be supervised by a physical educator or a physical therapist, and all programs include aerobic components (walking, cycling, dancing, jogging, swimming), muscle-strengthening activities (strength training, Pilates, functional exercise) and range of motion exercises (stretching, yoga, tai-chi).¹¹

Aerobic exercises increase the levels of peripheral beta-endorphins, which are correlated with decreased systemic sympathetic activity and improvement in serotonergic activity, reflected in the activity of neuromuscular junctions. Resistance training exercise promote better synchronization, recruitment and excitability of motor units. Finally, flexibility exercises can lead to a better control of articular structures and soft parts.^{38,39}

Now we describe particularities of physical activities and PE during treatment and follow-up periods of survivors after breast cancer treatment.

Physical exercises during breast cancer treatment

Breast cancer treatment should be individualized according to patients' age, hormonal status, comorbidities,

Table 1 – Description of physical activity intensities for the prevention and control of breast cancer

| Intensity | Description | Examples |
|--|--|---|
| Light <3 metabolic equivalents (METs) | Requires minimum effort, with small increase in HR and RR increase. On a 0-10 scale, perceived exertion ranges from 1 to 4. One can breathe calmly, talk, or even sing during exercise | Standing or sitting, washing dishes, doing arts and crafts |
| Moderate 3 – 5.9 METs | Requires greater physical effort, with perceptible but moderate increments in RR and HR. On a 0-10 scale, perceived exertion ranges from 5 to 6. One can talk with difficulty but not sing | Walking at > 5 Km/h; cycling at < 5 Km/h, double tennis, and ballroom dancing |
| Vigorous > 6 METs | Requires great physical effort with large increments in RR and HR. On a 0-10 scale, perceived exertion ranges from 7 to 8. One is not able to talk during the exercise. | Running, slope walking, bicycling at > 16 km/h, aerobic dancing |

Source: Adapted from Physical Activity Guidelines for the Brazilian Population.³⁷ HR: heart rate; RR: respiratory rate.

lifestyle, and personal choices, as well as to fundamental pillars in determining the prognosis: the extent of the disease (cancer staging) and the type of the tumor. In general, treatment can be divided into local therapy (surgery, radiotherapy, and breast reconstruction) and systemic therapy (chemotherapy, hormone therapy and biological therapy).⁴⁰

Chemotherapy is associated with fatigue, anorexia, anemia, neutropenia, thrombocytopenia, peripheral neuropathies and, in some cases, cardiotoxicity.^{41,42} The side effects of hormone therapy include weight gain, arthralgia, myalgia, bone loss, effects on the cardiovascular system and changes in the lipid profile.⁴³ Sequelae of radiation therapy include cardiac and pulmonary damage, lymphedema, brachial plexopathy and secondary malignant diseases.⁴⁴ Associated to these physical repercussions, emotional changes including depression, anxiety, low self-esteem, and negative body image can occur, since the disease affects an important symbol of femininity, sexuality and maternity.⁴⁵

Six months after the diagnosis, approximately 90% of women present at least one of the adverse symptoms of the antineoplastic therapy; 60% have multiple symptoms that affect not only patient therapy and quality of life, but also survival rates. Six years after treatment, up to 30% of women still have complaints related to the therapies.⁴⁶

Physical activity is safe and can be performed at different stages of cancer treatment, resulting in better quality of life, and improved global function, and lower psychological symptoms related to disease and its treatment.^{47,48}

Pain is one of the most common symptoms in breast cancer patients; 30-60% of patients have moderate to severe symptoms, that may lead to activity restrictions and

limitations in physical activity during and after therapeutical interventions.^{39,49} Pain manifestations tend to decrease with physical training, with direct implications for strength gain, better cardiorespiratory capacity and flexibility, and also for lower rates of fatigue, length of hospital stay, anxiety, depression, sleep disorders, nausea and vomiting.^{50,51} Figure 2 summarizes the final clinical effects of PE during treatment and in other breast cancer stages.

Van Waart et al.⁵² demonstrated that, in patients undergoing chemotherapy, an aerobic training program was associated with enhanced physical functioning, with maintenance of cardiorespiratory fitness, facilitating the return to work during and after treatment, in addition to reducing the incidence of nausea and vomiting as with the group that did not undergo physical training.⁵²

In addition, a multicentric study⁵³ evaluated the effect of physical activity in 301 patients during breast cancer chemotherapy and demonstrated improvements in health-related fitness in the three groups of exercise intervention: STAN group (three sessions of 25–30 min/week of vigorous-intensity aerobic exercise), HIGH group (three sessions of 150 min/week of vigorous-intensity aerobic exercise, and the COMB group (aerobic exercise plus a resistance exercise program). The higher dose of aerobic exercise intervention (HIGH group) was more effective in improving life quality and aerobic capacity, and in controlling pain and endocrine symptoms (e.g. hot flashes). However, the COMB group was superior to the HIGH and STAN groups in muscular strength gain.⁵³

In chemotherapy patient, resistance exercise programs are associated with improved self-esteem, muscular strength, and body composition, without causing or




Figure 2 – Clinical results of exercise in breast cancer timeline.

aggravating lymphedema or other complications in those who underwent surgery.^{54,55} Before initiating upper limbs exercises, it is recommended to assess the mobility of the arms. In addition, specific examination for the presence of peripheral neuropathies, musculoskeletal disorders, and risk of fractures, especially in patients on hormone therapy and in those with metastatic bone disease is advisable.^{46,56,57}

The combination of the three exercise programs proposed (aerobic, resistance and flexibility exercise) have largely contributed to the control of pain and fatigue. Improvement of cardiorespiratory function, due to increased aerobic capacity (maximal oxygen consumption) in combined exercise training programs may be explained by the ventilation perfusion matching and oxidative capacity of skeletal muscle.⁵⁵ This can play an important role in the management of structural disorder related to chemo- and radiation-induced toxicity.⁴⁶

Chemotherapeutic agents that also cause direct and indirect cardiotoxicity, with acceleration of general and vascular aging, and consequent decline of cardiopulmonary reserve. Both the disease and the therapy can contribute to weight gain and reduction of physical activity,⁵⁸ potentially increasing the risk for cardiovascular diseases (CVD). Studies on secondary prevention have corroborated the improvement of cardiopulmonary function with physical training programs in breast cancer women.⁵⁹

Cardiotoxicity, associated with psychoemotional factors, affects autonomic balance and consequently cardiovascular mortality.⁶⁰ In patients treated at initial stages of the disease, a sustained increment in sympathetic function and reduction in parasympathetic effect on the sinoatrial node have been reported.⁶¹ Other reports have pointed out a reduction in heart rate variability and baroreflex sensitivity among women with a history of breast cancer.⁶¹⁻⁶³

In addition to autonomic regulation, another important factor for the development of CVD is endothelial dysfunction. A recent meta-analysis⁶⁴ evaluated 163 patients from four studies (two on breast cancer, two on prostate cancer). Aerobic exercise improved vascular function and peak oxygen consumption. These data reinforce the importance of PE as an adjuvant therapy in breast cancer treatment, especially regarding the management of the side effects.

With respect to overall and specific mortality, evidence accumulated so far suggests a favorable effect of moderate to vigorous physical activity, and preliminary evidence indicates associations of physical activity with risk reduction of breast cancer recurrence and progression.¹¹ A recent study evaluated systematic reviews on physical activity and reduction of all-cause mortality and breast cancer mortality, considering the dose-response relationship (also regardless of body mass index); the certainty of evidence was moderate. Regarding associations of the domain or the type of physical activity with mortality, the certainty of evidence was classified as low and, so far, it was not possible to identify physical activity modalities that had the greatest impact on the outcome. The same study evaluated risks and benefits, patients' values and preferences,

resources required to meet the recommended physical activity, equity, acceptability of the recommendation, and the strength of the recommendation of PE to increase breast cancer survival was classified as "strong".¹¹ Future studies may change the quality of available evidence; it remains an open field of research.

Physical exercises in breast cancer post-treatment

PE have been strongly recommended for breast cancer survivors and associated not only with improvement of quality of life, but also with possible increased survival.⁶⁵⁻⁶⁷ A prospective study that included 2987 women with stage I, II, or III breast cancer between 1984 and 1988, followed up until death or June 2002, showed that physical activity after breast cancer diagnosis can reduce the risk of death for the disease. The greatest benefit occurred in women who performed the equivalent of walking three to five hours per week at an average pace.⁶⁵

After the end of treatment, the main objective is to rehabilitate patients to return to their usual activities. Regular exercises can contribute to physical and psychological well-being and improved quality of life, consisting of one of the main recommendations to prevent chronic degenerative conditions; it would not be different for patients who had recently faced cancer treatment.⁶⁸ Evidence has shown that PE have a positive impact on survival and minimize breast cancer-related morbidity.⁶⁵ Despite favorable data, the practice of PE is limited by barriers like fatigue, lack of motivation, loss of self-confidence, inadequate follow-up, lack of family support and lack of instructions.

Encouraging women in the post-treatment period to adopt a healthy lifestyle – by avoiding excess alcohol and increasing fruit and vegetable intake and physical activity volume, is important to improve their quality of life and health.⁶⁹ Increased exercise levels represent a modifiable health behavior that can ameliorate sequelae of the disease and help women to return to their health status before cancer diagnosis and treatment.⁷⁰ Thus, current recommendations of PE for breast cancer survivors are based on the return, as soon as possible, to habitual daily activities, on the maintenance of metabolic expenditure during and after therapies, and on the classical recommendation of weekly aerobic exercises.⁵⁷

Also, it has been demonstrated that physical inactivity is related to weight gain after the diagnosis which, in turn, has been associated with lower survival in some studies.^{71,72} More physically active women are less likely to gain weight after the diagnosis, improving the chance of survival.^{65,73}

Obesity is related to increased mortality rates for breast cancer (13-20%) and all-cause mortality (14-70%).⁷⁴⁻⁷⁷ Obesity was also associated with a twice greater chance of post-menopausal contralateral breast cancer and a nearly 60% greater occurrence of other cancers.⁷⁶ Therefore, maintaining normal body mass index can reduce the risk of a new breast cancer in the postmenopausal period, other cancers, and all-cause mortality.^{75,76,78}

Giallauria et al.⁶¹ evaluated whether exercise training improved autonomic function in women with a history of primary invasive breast cancer. Fifty-one patients enrolled in the DIANA clinical trial were divided into two groups. Group 1 (n = 25) that followed a formal exercise program of moderate intensity – three sessions/week on a bicycle or treadmill at $70 \pm 2\%$ VO_2 peak for 12 weeks, followed by one session/week until one-year follow-up. Group 2 while a control group (n = 26) did not perform any formal training. At baseline and at one-year follow-up, all patients underwent cardiopulmonary exercise stress test. Heart rate recovery (HRR) was calculated as the difference between heart rate at peak exercise and heart rate at the first minute of recovery. Compared with control group, group 1 showed significant improvement of VO_2 peak (from 12.4 ± 2.9 to 14.3 ± 3.3 mL/kg/min; $p < 0.001$) and in HRR (from 17.6 ± 6.4 to 23.0 ± 8.3 beats/min; $p < 0.001$). The authors concluded that moderate-intensity exercise is associated with improvement of autonomic function in breast cancer survivors.⁶²

Evidence suggests that PE can also promote physiological and psychological benefits in cancer survivors.^{70,79} A meta-analysis of randomized controlled trials by Fong et al.⁷⁰ reported that PE had positive effects on physical functions, body weight and quality of life in patients after treatment for breast cancer.⁷⁰ Additionally, results from another systematic review⁶⁶ indicated that PE can have beneficial effects on overall and on certain domains of quality of life, like body image, self-esteem, emotional well-being, sexuality, sleep disorder, social functioning, anxiety, fatigue and pain.⁶⁶ Also, a Cochrane database systematic review that included 63 trials and 5761 women evaluated the effects of PE in patients in the post-treatment period and in a control group. Once again, it was shown that physical activity interventions resulted in improvements in quality of life, emotional health, anxiety, physical function, muscular strength, and fatigue. Besides, relatively few adverse events were reported in the trials, suggesting that PE are safe in this population.⁶⁷

To be safe, exercise prescription requires the understanding, by the multidisciplinary team (physical educators, physical therapists, among others), of the peculiarities, implications and consequences of cancer treatment.⁵⁷ Prescriptions should be made according to pre-treatment physical performance and comorbidities of cancer survivors, therapeutic response and negative immediate and persistent effects of treatment.⁵⁷ Special attention should be given to peripheral neuropathies and secondary musculoskeletal diseases, regardless of the treatment time. Patients on hormone therapy should be assessed for the risk of fractures. It is also recommended evaluating the mobility of arms and shoulders before initiating upper limb exercises. It is important to consider the needed time for wound healing, which may be eight weeks or longer in mastectomies.⁵⁷

Individuals with bone metastatic disease will required individualized exercise programming aiming at determining safety limits before initiating the PE. Rehabilitation of these patients include adaptations in pre-established programs,

with reductions in impact, intensity, and volume, due to the increased risk posed by bone frailty and fractures.⁵⁷ In addition, individuals with known CVD (secondary to cancer or not) also require initial individualized examination regarding the safety of the exercise programs, closer supervision, and shorter intervals. Guidelines' recommendations on exercise and rehabilitation should be followed, especially considering cardiovascular and pulmonary contraindications.⁵⁷

Cancer survivors should be physically active. However, exercise prescription including exercise frequency, intensity, type and duration has been based on limited literature data. Table 2 summarized recommended exercise prescription for breast cancer patients in the post-treatment.^{47,57}

The progression of PE should be slower in cancer survivors as compared with healthy individuals, particularly if the prescribed exercises result in greater fatigue and unexpected adverse effects, which serves as an alert to individual's capacity thresholds. There are no maximum loads for weight training exercises for these patients. Attention should be paid to symptoms in the arms and shoulders, including lymphedema, resulting in load decrease or interruption of specific exercises according to the symptom reported.⁵⁷

Despite all the benefits of regular physical activity described above, there is no consensus or clear standard of the magnitude of their benefits, the way of administration or the most effective PE for this population. Further research is needed to establish the ideal exercise prescription. Studies so far have evaluated the effects of different exercise modalities, frequencies, intensities and durations on specific outcomes in breast cancer survivors, which make generalization and standardization of results difficult.

Finally, we reinforce the need of all health professionals involved – physicians, physical educators, physical therapists, psychologists, and nutritionists – be aware of the importance of encouraging these women to regularly exercise after breast cancer treatment, highlighting the benefits and excellent cost-effectiveness.

Conclusion

Regular exercise/physical activity should be encouraged among women, targeting primary prevention, improvement of life quality and reduction of mortality among survivors, although studies have not reported the strength of evidence for breast cancer control. It is also important to highlight the important role of PE in reducing the incidence of CVD, which reinforces the importance of encouraging these women to be physically active. Although attention should be paid to some details in exercise prescription to breast cancer patients, in general, it is not very different from that made to the general population. Future studies are needed to better guide individualized prescriptions for these patients.

Table 2 – Recommended prescription of physical exercises in the post-treatment of breast cancer

| | Frequency | Intensity | Duration/performance | Quality |
|-------------|---|---|---|---|
| Aerobic | Patient should start exercises twice a week and progressively increase until 3-5 times a week | Patient should be instructed about effort perception. If exercises are well tolerated (without symptoms or side effects), the intensity of exercises should not be different from that in healthy population. Intensity should be from moderate to vigorous | Exercise duration should be increased according to patient tolerance. A target of 75min/week of vigorous exercise or 150min/week of moderate exercise should be aimed | Rhythmic, prolonged exercises that use large muscle groups. Examples: swimming, walking, bicycling, dancing |
| Resistance | 2-3 days a week | Moderate intensity (60-70% maximum repetitions) | Sets of 8 - 12 repetitions | Programs should include weights, resistance training equipment, functional tasks with weights, using the main large muscle groups |
| Flexibility | Exercises can be performed daily, according to patient's condition | | | Programs should include stretching exercises. Attention to body parts with restricted mobility due to treatment |

Source: Adapted from the American College of Sports Medicine guidelines for exercise testing and prescription.⁴⁷

Author Contributions

Conception and design of the research: Campos MSB, Feitosa RHF, Mizzaci CC, Flach MRTV, Siqueira BJM; Acquisition of data: Campos MSB, Feitosa RHF, Mizzaci CC, Flach MRTV, Siqueira BJM; Analysis and interpretation of the data, Writing of the manuscript and Critical revision of the manuscript for important intellectual content: Campos MSB, Feitosa RHF, Mizzaci CC, Flach MRTV, Siqueira BJM, Mastrocolla LE.

Potential Conflict of Interest

No potential conflict of interest relevant to this article was reported.

Sources of Funding

There were no external funding sources for this study.

Study Association

This study is not associated with any thesis or dissertation work.

References

- Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, Bray F. Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. *CA Cancer J Clin*. 2021;71(3):209-49. doi:10.3322/caac.21660
- Instituto Nacional de Câncer José Alencar Gomes da Silva. Estimativa | 2020 Incidência de Câncer no Brasil. Rio de Janeiro; 2019. v.1;p:1-122. ISBN:978-85-7318-389-4 [Acesso em 26 jul, 2022] Disponível em: <https://www.inca.gov.br/estimativa/taxas-ajustadas/neoplasia-maligna-da-mama-feminina-e-colo-do-utero>.
- Wilkinson L, Gathani T. Understanding breast cancer as a global health concern. *Br J Radiol*. 2022;95(1130):20211033. doi:10.1259/bjr.20211033
- Gonçalves, A, Jobim, P, Vanacor R, Nunes L, Albuquerque IM, Bozzetti MC, et al Câncer de mama: mortalidade crescente na Região Sul do Brasil entre 1980 e 2002. *Cad. Saúde Pública*. 2007; 23 (8) <https://doi.org/10.1590/S0102>.
- Instituto Nacional de Câncer José Alencar Gomes da Silva. In: Atlas On-line de Mortalidade. Rio de Janeiro: INCA, 2022. Ministério da Saúde. [Acesso em 26 jul.2022] Disponível em: <https://www.inca.gov.br/MortalidadeWeb/pages/Modelo01/consultar.xhtml>
- Collaborative Group on Hormonal Factors in Breast Cancer. Familial breast cancer: collaborative reanalysis of individual data from 52 epidemiological studies including 58,209 women with breast cancer and 101,986 women without the disease. *Lancet*. 2001;358(9291):1389-99. doi:10.1016/S0140-6736(01)06524-2
- National Breast Cancer Foundation, Inc. [Internet] [Cited in 2022 Mar 07] Available from: <https://www.nationalbreastcancer.org>; <https://www.cancer.gov/about-cancer/causes-prevention/genetics/brca-fact-sheet>, downloaded
- Castelló A, Martín M, Ruiz A, Casas AM, Baena-Canada JM, Lope V, et al. Lower Breast Cancer Risk among Women following the World Cancer Research Fund and American Institute for Cancer Research Lifestyle Recommendations: EpiGEICAM Case-Control Study. *PLoS One*. 2015;10(5):e0126096. doi:10.1371/journal.pone.0126096
- Ashcraft KA, Peace RM, Betof AS, Dewhirst MW, Jones LW. Efficacy and Mechanisms of Aerobic Exercise on Cancer Initiation, Progression, and Metastasis: A Critical Systematic Review of In Vivo Preclinical Data. *Cancer Res*. 2016;76(14):4032-50. doi:10.1158/0008-5472.CAN-16-0887
- World Cancer Research Fund/American Institute for Cancer Research. Diet, Nutrition, Physical Activity and Cancer: a Global Perspective. Continuous Update Project Expert Report 2018. the Cancer Process. [Cited in 2022 Apr Available from: <https://docslib.org/doc/3676076/continuous-update-project-expert-report.2018>

11. Sociedade Brasileira de Oncologia Clínica(SBOC).Sociedade Brasileira de Atividade Física e Saúde. Instituto Nacional de Cancer José de Alencar Gomes da Silva (INCA), Atividade Física e Câncer: Recomendações Para Prevenção e Controle. São Paulo, 2022, v. 01, p. 1-58.
12. Pedersen L, Christensen JF, Hojman P. Effects of exercise on tumor physiology and metabolism. *Cancer J*. 2015;21(2):111-6. doi:10.1097/PPO.0000000000000096
13. McTiernan A. Mechanisms linking physical activity with cancer. *Nat Rev Cancer*. 2008;8(3):205-11. doi:10.1038/nrc2325
14. Betof AS, Lascola CD, Weitzel D, London C, Scarbrough PM, Gayathri R, et al. Modulation of murine breast tumor vascularity, hypoxia and chemotherapeutic response by exercise. *J Natl Cancer Inst*. 2015;107(5):djv040. doi:10.1093/jnci/djv040.
15. Ruiz-Casado A, Martín-Ruiz A, Pérez LM, Provencio M, Fiuza-Luces C, Lucia A. Exercise and the Hallmarks of Cancer. *Trends Cancer*. 2017;3(6):423-41. doi:10.1016/j.trecan.2017.04.007
16. Hojman P. Exercise protects from cancer through regulation of immune function and inflammation. *Biochem Soc Trans*. 2017;45(4):905-11. doi:10.1042/BST20160466
17. Onuchic AC, Chammas R. Câncer e o microambiente tumoral. *Rev Med*. 2010;89(1):21-31. DOI: 10.11606/issn.1679-9836.v89i1p21-31.
18. Suarez-Carmona M, Lesage J, Cataldo D, Gilles C. EMT and inflammation: inseparable actors of cancer progression. *Mol Oncol*. 2017;11(7):805-23. doi:10.1002/1878-0261.12095.
19. Hanahan D, Weinberg RA. Hallmarks of cancer: the next generation. *Cell*. 2011;144(5):646-74. doi:10.1016/j.cell.2011.02.013
20. Dethlefsen C, Pedersen KS, Hojman P. Every exercise bout matters: linking systemic exercise responses to breast cancer control. *Breast Cancer Res Treat*. 2017;162(3):399-408. doi:10.1007/s10549-017-41294
21. Hojman P, Gehl J, Christensen JF, Pedersen BK. Molecular Mechanisms Linking Exercise to Cancer Prevention and Treatment. *Cell Metab*. 2018;27(1):10-21. doi:10.1016/j.cmet.2017.09.015
22. Pedersen L, Idorn M, Olofsson GH, Lauenborg B, Nookaew I, Hansen RH, et al. Voluntary Running Suppresses Tumor Growth through Epinephrine- and IL-6-Dependent NK Cell Mobilization and Redistribution. *Cell Metab*. 2016;23(3):554-62. doi:10.1016/j.cmet.2016.01.011.
23. Rundqvist H, Veliça P, Barbieri L, Gameiro PA, Bargiela D, Gokjovic M, et al. Cytotoxic T-cells mediate exercise-induced reductions in tumor growth. *Elife*. 2020;9:e59996. doi:10.7554/eLife.59996
24. Toffoli EC, Sweegers MG, Bontkes HJ, Verheul HMW, vander Vliet HJ, Altenburg TM, et al. Effects of physical exercise on natural killer cell activity during (neo)adjuvant chemotherapy: A randomized pilot study. *Physiol Rep*. 2021;9(11):e14919. doi:10.14814/phy2.14919
25. Alizadeh AM, Isanejad A, Sadighi S, Mardani M, Kalaghchi B, Hassan ZM. High-intensity interval training can modulate the systemic inflammation and HSP70 in the breast cancer: a randomized control trial. *J Cancer Res Clin Oncol*. 2019;145(10):2583-93. doi:10.1007/s00432-019-02996-y
26. Rockhill B, Willett WC, Hunter DJ, Manson JE, Hankinson SE, Colditz GA. A prospective study of recreational physical activity and breast cancer risk. *Arch Intern Med*. 1999;159(19):2290-6. doi:10.1001/archinte.159.19.2290
27. McTiernan A, Kooperberg C, White E, Coates R, Adams-Campbell LL, Woods N, et al. Recreational physical activity and the risk of breast cancer in postmenopausal women: the Women's Health Initiative Cohort Study. *JAMA*. 2003;290(10):1331-6. doi:10.1001/jama.290.10.1331
28. Friedenreich CM, Cust AE. Physical activity and breast cancer risk: impact of timing, type and dose of activity and population subgroup effects. *Br J Sports Med*. 2008;42(8):636-47. doi:10.1136/bjsm.2006.029132
29. Lynch BM, Neilson HK, Friedenreich CM. Physical activity and breast cancer prevention. *Recent Results Cancer Res*. 2011;186:13-42. doi:10.1007/978-3-642-04231-7_2
30. Friebe TM, Domchek SM, Rebbeck TR. Modifiers of cancer risk in BRCA1 and BRCA2 mutation carriers: systematic review and meta-analysis [published correction appears in *J Natl Cancer Inst*. 2014 Aug;106(8):dju235. doi:10.1093/jnci/dju235. *J Natl Cancer Inst*. 2014;106(6):dju091.
31. Wu Y, Zhang D, Kang S. Physical activity and risk of breast cancer: a meta-analysis of prospective studies. *Breast Cancer Res Treat*. 2013;137(3):869-82. doi:10.1007/s10549-012-2396-7
32. Brown JC, Winters-Stone K, Lee A, Schmitz KH. Cancer, physical activity, and exercise. *Compr Physiol*. 2012;2(4):2775-809. doi:10.1002/cphy.c120005
33. Rezende LFM, Sá TH, Markozannes G, Rey-López JP, Lee I-M, Konstantinos K, Tsilidis KK, et al. Physical activity and cancer: an umbrella review of the literature including 22 major anatomical sites and 770 000 cancer cases. *Br J Sports Med*. 2018;52(13):826-33. doi:10.1136/bjsports-2017-098391
34. Effects of exercise training on fasting insulin, insulin resistance, insulin-like growth factors, and insulin-like growth factor binding proteins in postmenopausal breast cancer survivors: a randomized controlled trial. *Cancer Epidemiol Biomarkers Prev*. 2003;12(8):721-7. PMID: 12917202
35. de Boer MC, Wörner EA, Verlaan D, van Leeuwen PAM. The Mechanisms and Effects of Physical Activity on Breast Cancer. *Clin Breast Cancer*. 2017;17(4):272-8. doi:10.1016/j.clbc.2017.01.006
36. World Health Organization.(WHO). Guidelines on physical activity and sedentary behaviour: web annex: evidence profiles. World Health Organization. <https://apps.who.int/iris/handle/10665/336657>. Licença: CC BY-NC-SA 3.0 IGO
37. Brasil. Ministério da Saúde. Secretaria de Atenção Primária à Saúde, Departamento de Promoção da Saúde. Brasília ;2021. 54p. [Internet] [Citado em 2022 maio 14] Disponível em: http://bvsm.sau.gov.br/bvs/publicacoes/guia_atividade_fisica_populacao_brasileira.pdf ISBN978-85-334-2885-0
38. McArdle WD, Katch FI, Katch VL. Exercise physiology: energy, nutrition and human performance. 2nd ed. Rio de Janeiro: Guanabara Koogan; 2011. 1099p.
39. Reis AD, Pereira PTV, Diniz RR, 58Castro Filha JG, Santos A, Ramallo BT, et al. Effect of exercise on pain and functional capacity in breast cancer patients. *Health Qual Life Outcomes*. 2018;16(1):58. DOI: 10.1186/s12955-018-0882-2
40. Soerjomataram I, Louwman MW, Ribot JC, Roukema JA, Coebergh JW. An overview of prognostic factors for long-term survivors of breast cancer. *Breast Cancer Res Treat*. 2008 Feb;107(3):309-30. doi: 10.1007/s10549-007-9556-1.
41. Ramírez K, Acevedo F, Herrera ME, Ibáñez C, Sánchez C. Actividad física y cáncer de mama: un tratamiento dirigido [Physical activity and breast cancer]. *Rev Med Chil*. 2017;145(1):75-84. doi:10.4067/S0034-98872017000100011
42. Barry E, Alvarez JA, Scully RE, Miller TL, Lipshultz SE. Anthracycline-induced cardiotoxicity: course, pathophysiology, prevention and management. *Expert Opin Pharmacother*. 2007;8(8):1039-58. doi:10.1517/14656566.8.8.1039
43. Conte P, Frassoldati A. Aromatase inhibitors in the adjuvant treatment of postmenopausal women with early breast cancer: Putting safety issues into perspective. *Breast J*. 2007;13(1):28-35. doi:10.1111/j.1524-4741.2006.00359.x
44. Senkus-Konefka E, Jassem J. Complications of breast-cancer radiotherapy. *Clin Oncol (R Coll Radiol)*. 2006;18(3):229-35. doi:10.1016/j.clon.2005.11.004
45. Groenvold M, Petersen MA, Idler E, Björner JB, Fayes PM, Mouridsen HT. Psychological distress and fatigue predicted recurrence and survival in primary breast cancer patients. *Breast Cancer Res Treat*. 2007;105(2):209-19. doi:10.1007/s10549-006-9447-x
46. Schmitz KH, Speck RM, Rye SA, DiSipio T, Hayes SC. Prevalence of breast cancer treatment sequelae over 6 years of follow-up: the Pulling Through Study. *Cancer*. 2012;118(8 Suppl):2217-25. doi:10.1002/cncr.27474

47. Diretrizes do ACSM para os Testes de Esforço e sua Prescrição. 10. ed., São Paulo: Guanabara Koogan; 2019.v.1.p:314-25. ISBN 9788527732871.
48. Bouillet T, Bigard X, Bami C, Chouahnia C, Copel L, Dauchy, S, et al. Role of physical activity and sport in oncology: scientific commission of the National Federation Sport and Cancer CAMI. *Crit Rev Oncol Hematol*. 2015;94(1):74-86. doi:10.1016/j.critrevonc.2014.12.012
49. Garcia LB, Guirro EC, Efeitos da estimulação de alta voltagem no linfedema pós-mastectomia. *Rev Bras Fisioter*. 2005;9(2):243-8.
50. Yates P, Bashford J, Pyke C, et al. Hayes SC, Rye S, Disipio T, et al. Exercise for health: a randomized, controlled trial evaluating the impact of a pragmatic, translational exercise intervention on the quality of life, function and treatment-related side effects following breast cancer. *Breast Cancer Res Treat*. 2013;137:175-86. Doi: 10.1007/s10549-012-2331-y
51. Buffart LM, Sweegers MC, May A, Chinapaw MJ, van Vulpen JK, Rob U Newton RU, et al. Targeting Exercise Intervention]s to Patients With Cancer in Need: An Individual Patient Data Meta-Analysis. *J Natl Cancer Inst*. 2018;110(11):1190-200. doi:10.1093/jnci/djy161
52. van Waart H, Stuiver MM, van Harten WH, Geleijn L, Kieffer JM, Buffart LM, et al. Effect of Low-Intensity Physical Activity and Moderate- to High-Intensity Physical Exercise During Adjuvant Chemotherapy on Physical Fitness, Fatigue, and Chemotherapy Completion Rates: Results of the PACES Randomized Clinical Trial. *J Clin Oncol*. 2015;33(17):1918-27. doi:10.1200/JCO.2014.59.1081.
53. Courneya KS, McKenzie DC, Mackey JR, Gelmon K, Friedenreich CM, Yasui Y, et al. Effects of exercise dose and type during breast cancer chemotherapy: multicenter randomized trial. *J Natl Cancer Inst*. 2013;105(23):1821-1832. doi:10.1093/jnci/djt297
54. Courneya K, Lee J, Fairey A, Campbell K, Ladha A, Friedenreich C, et al. Physical activity in cancer survivors: implications for recurrence and mortality. *Cancer Ther* 2004; 2: 1-12.
55. Sauer DP, Perez JA, Carletti L. Effect of training on ventilatory efficiency in healthy subjects. *Rev Bras Med Esp*. 2014;20(6):470-3. <https://doi.org/10.1590/1517-86922014200601814>.
56. Schneider CM, Dennehy CA, Roozeboom M, Carter SD. A model program: exercise intervention for cancer rehabilitation. *Integr Cancer Ther*. 2002;1(1):76-82. doi:10.1177/153473540200100117
57. Schmitz KH, Courneya KS, Mathews C, Demark-Wahnefried W, Galvão D, Pinto B, et al. American College of Sports Medicine roundtable on exercise guidelines for cancer survivors. *Med Sci Sports Exerc*. 2010;42(7):1409-26. doi:10.1249/MSS.0b013e3181e0c112
58. Rock, CL, Flatt SW, Newman V, Caan BJ, Haan MN, Stefanick ML, et al. Factors associated with weight gain in women after diagnosis of breast cancer. Women's Healthy Eating and Living Study Group. *J Am Diet Assoc*. 1999;99(10):1212-21. doi:10.1016/s0002-8223(99)00298-9
59. Travier N, Velthuis M, Bisschop C, van den Beys B, Monnikhof EM, Los M, et al. Effects of an 18-week exercise programme started early during breast cancer treatment: a randomised controlled trial. *BMC Med*. Doi:10.1186/s12916-015-0362-z
60. Fagundes CP, Murray DM, Hwang BS, Gouin JP, Thayer JF, Sollers 3rd JJ, et al. Sympathetic and parasympathetic activity in cancer-related fatigue: more evidence for a physiological substrate in cancer survivors. *Psychoneuroendocrinology*. 2011;36(8):1137-47. doi:10.1016/j.psyneuen.2011.02.005
61. Giallauria F, Vitelli A, Maresca L, et al. Exercise training improves cardiopulmonary and endothelial function in women with breast cancer: findings from the Diana-5 dietary intervention study. *Intern Emerg Med*. 2016;11(2):183-189. doi:10.1007/s11739-015-1259-8.
62. Giallauria F, Maresca L, Vitelli A, Santucci de Magistris M, Chiodini P, Mattiello A, Gentile M, et al. Exercise training improves heart rate recovery in women with breast cancer. *Springerplus*. 2015;4:388. doi:10.1186/s40064-015-1179-0
63. Spence RR, Heesch KC, Brown WJ. Exercise and cancer rehabilitation: a systematic review. *Cancer Treat Rev*. 2010;36(2):185-94. doi:10.1016/j.ctrv.2009.11.003
64. Beaudry RI, Liang Y, Boyton ST, Tucker W, Brothers MR, Daniel KM, et al. Meta-analysis of Exercise Training on Vascular Endothelial Function in Cancer Survivors. *Integr Cancer Ther*. 2018;17(2):192-9. doi:10.1177/1534735418756193
65. Holmes MD, Chen WY, Feskanich D, Kroenke CH, Colditz GA. Physical activity and survival after breast cancer diagnosis. *JAMA*. 2005;293(20):2479-86. doi:10.1001/jama.293.20.2479
66. Mishra SI, Scherer RW, Geigle PM, Berlanstein DR, Topaloglu O, Gotay CC, Snyder C, et al. Exercise interventions on health-related quality of life for cancer survivors. *Cochrane Database Syst Rev*. 2012;2012(8):CD007566. doi:10.1002/14651858.CD007566.pub2
67. Lahart IM, Metsios GS, Nevill AM, Carmichael AR. Physical activity for women with breast cancer after adjuvant therapy. *Cochrane Database Syst Rev*. 2018;1(1):CD011292. doi:10.1002/14651858.CD011292.pub2
68. Marshall SJ, Jones DA, Ainsworth BE, Reis JP, Levy SS, Macera CA. Race/ethnicity, social class, and leisure-time physical inactivity. *Med Sci Sports Exerc*. 2007;39(1):44-51. doi:10.1249/01.mss.0000239401.16381.37.
69. Demark-Wahnefried W, Aziz NM, Rowland JH, Pinto BM. Riding the crest of the teachable moment: promoting long-term health after the diagnosis of cancer. *J Clin Oncol*. 2005;23(24):5814-30. doi:10.1200/JCO.2005.01.230
70. Fong DY, Ho JW, Hui BP, Lee AM, McFarlane D, Leung SSK, et al. Physical activity for cancer survivors: meta-analysis of randomised controlled trials. *BMJ*. 2012;344:e70. doi:10.1136/bmj.e70
71. Camoriano JK, Loprinzi CL, Ingle JN, Therneau TM, Krook JE, Veeder MH. Weight change in women treated with adjuvant therapy or observed following mastectomy for node-positive breast cancer. *J Clin Oncol*. 1990;8(8):1327-34. doi:10.1200/JCO.1990.8.8.1327
72. Kroenke CH, Chen WY, Rosner B, Holmes MD. Weight, weight gain, and survival after breast cancer diagnosis. *J Clin Oncol*. 2005;23(7):1370-8. doi:10.1200/JCO.2005.01.079.
73. Lahmann PH, Schulz M, Hoffmann K, Boeing H, Tonnelland A, Olsen A, et al. Long-term weight change and breast cancer risk: the European prospective investigation into cancer and nutrition (EPIC). *Br J Cancer*. 2005;93(5):582-9. doi:10.1038/sj.bjc.6602763
74. Berclaz G, Li S, Price KN, Coates AS, Castiglione-Gertsch M, Rudenstam C-M, et al. Body mass index as a prognostic feature in operable breast cancer: the International Breast Cancer Study Group experience. *Ann Oncol*. 2004;15(6):875-84. doi:10.1093/annonc/mdh222
75. Dignam JJ, Wieand K, Johnson KA, Fisher B, Xu L, Mamounas EP. Obesity, tamoxifen use, and outcomes in women with estrogen receptor-positive early-stage breast cancer. *J Natl Cancer Inst*. 2003;95(19):1467-76. doi:10.1093/jnci/djg060
76. Dignam JJ, Wieand K, Johnson KA, Raich P, Anderson SJ, Somkin C, et al. Effects of obesity and race on prognosis in lymph node-negative, estrogen receptor-negative breast cancer. *Breast Cancer Res Treat*. 2006;97(3):245-54. doi:10.1007/s10549-005-9118-3
77. Holmes MD, Stampfer MJ, Colditz GA, Rosner B, Hunter DJ, Willett WC. Dietary factors and the survival of women with breast carcinoma. *Cancer*. 1999;86(5):826-35. doi:10.1002/(sici)1097-0142(19990901)86:5<826::aid-cnrcr19>3.0.co;2-0
78. Byers T, Sedjo RL. A weight loss trial for breast cancer recurrence: pre-menopausal, post-menopausal, both, or neither? *Cancer Causes Control*. 2006;17(1):1-3. doi:10.1007/s10552-005-0547-4
79. Brown JC, Huedo-Medina TB, Pescatello LS, Ryan SM, Pescatello SM, Moker E, et al. The efficacy of exercise in reducing depressive symptoms among cancer survivors: a meta-analysis. *PLoS One*. 2012;7(1):e309255 doi:10.1371/journal.pone.0030955



This is an open-access article distributed under the terms of the Creative Commons Attribution License