# **Short Editorial**



## Understanding the Link between Visceral Fat and Heart Health

Nuno Bettencourt<sup>1</sup>

Universidade do Porto Faculdade de Medicina - Unidade de Investigação Cardiovascular,¹ Porto – Portugal Short Editorial related to the article: Resting Heart Rate Variability is Independently Associated with Visceral Fat Rating Scores in Saudi Adult Males

Cardiovascular disease (CVD) remains a significant global health concern, with obesity and its associated metabolic abnormalities contributing substantially to CVD risk. In recent years, research has increasingly focused on the role of visceral adipose tissue (VAT) in cardiovascular health, recognizing its unique contribution to metabolic dysfunction and cardiovascular risk.<sup>1,2</sup>

Heart rate variability (HRV) is a measure of the variation in the time intervals between consecutive heartbeats, reflecting the balance between the sympathetic and parasympathetic nervous systems – and, therefore, serving as a valuable indicator of autonomic function and cardiovascular health. Reduced HRV is associated with increased cardiovascular risk, making it a potentially useful tool for risk assessment and monitoring in different clinical scenarios.<sup>3-5</sup>

In this number of the journal, Habib SS et al., studied the association between VAT, as measured by visceral fat rating (VFR), and HRV in a cohort of 99 Saudi healthy men.<sup>6</sup> The purpose of the investigation was to test the hypothesis that HRV differs between participants as a function of VAT categorization, and that indexes of VAT are more strongly associated with HRV than relative fat mass in healthy adult men.

Despite some important limitations, namely the cross-sectional design (limiting the ability to establish causality) and sample composition of healthy men (limiting the generalizability to other populations), this interesting study offers valuable insights into the complex interplay between adiposity and autonomic nervous system function and may be of clinical relevance for risk-assessment and primary prevention guidance.

One of the most striking findings of the study was the significant association between VFR and HRV parameters, particularly "root-mean-square of successive differences"

(RMSSD) and "standard deviation of normal RR intervals" (SDNN), which are indicative of parasympathetic modulation and global variability, respectively. Interestingly, individuals with higher levels of visceral fat exhibited reduced HRV, suggesting impaired autonomic function. Importantly, these associations remained significant even after adjusting for potential confounders such as age, BMI, and blood pressure, underscoring the robustness of the findings.

Moreover, the study highlights the sensitivity of HRV parameters to changes in visceral fat, surpassing traditional measures such as body fat percentage and muscle mass to visceral fat ratio (MMVFR). This underscores the potential utility of HRV metrics as sensitive markers for monitoring cardiac-autonomic status in response to interventions targeting visceral fat reduction.

This study represents a significant step forward in our understanding of the relationship between visceral fat and heart health. By identifying visceral fat as a modifiable risk factor for impaired autonomic function, the findings underscore the importance of targeted interventions aimed at reducing visceral fat accumulation to improve cardiovascular health. Lifestyle modifications aimed at reducing visceral fat, such as dietary changes and increased physical activity, could potentially improve autonomic function, and mitigate cardiovascular risk.

Furthermore, this study underscores the importance of ongoing research in this area to elucidate the underlying mechanisms linking visceral fat accumulation to autonomic dysfunction. As new perspectives for the assessment of cardiovascular risk and tailored interventions are clearly open with these lines of research, future studies may explore the effectiveness of interventions targeting visceral fat reduction in improving HRV and cardiovascular outcomes.

#### **Keywords**

Heart Rate Variability; Visceral Adipose Tissue

Mailing Address: Nuno Bettencourt •

Universidade do Porto Faculdade de Medicina - Unidade de Investigação Cardiovascular - Alameda Professor Hernâni Monteiro, 4200-319 Porto - Portugal E-mail: bettencourt.n@gmail.com

Manuscript received April 29, 2024, revised manuscript May 08, 2024, accepted May 08, 2024

DOI: https://doi.org/10.36660/abc.20240288i

### **Short Editorial**

#### References

- Guimarães GC Filho, Silva LT, Silva RMCE. Correlation Among Waist Circumference and Central Measures of Blood Pressure. Arq Bras Cardiol. 2022;119(2):257-64. doi: 10.36660/abc.20210432.
- Santos CPC, Lagares LS, Santos SRM, Silva MSP, Macedo RC, Almeida LAB, et al. Association Between Arterial Hypertension and Laboratory Markers, Body Composition, Obstructive Sleep Apnea and Autonomic Parameters in Obese Patients. Arq Bras Cardiol. 2023;120(7):e20220728. doi: 10.36660/ abc.20220728.
- Mota GAF, Gatto M, Pagan LU, Tanni SE, Okoshi MP. Diabetes Mellitus, Physical Exercise and Heart Rate Variability. Arq Bras Cardiol. 2023;120(1):e20220902. doi: 10.36660/abc.20220902.
- Gomes BFO, Benchimol-Barbosa PR, Nadal J. Predictive Model of All-Cause Death in Patients with Heart Failure using Heart Rate Variability. Arq Bras Cardiol. 2023;120(11):e20220379. doi: 10.36660/abc.20220379.
- Souza EG Neto, Peixoto JVC, Rank Filho C, Petterle RR, Fogaça RTH, Wolska BM, et al. Effects of High-Intensity Interval Training and Continuous Training on Exercise Capacity, Heart Rate Variability and Isolated Hearts in Diabetic Rats. Arq Bras Cardiol. 2023;120(1):e20220396. doi: 10.36660/abc.20220396.
- Habib SS, Alkahtani S, Aljawini N, Habib SM, Flatt AA. A Variabilidade da Frequência Cardíaca em Repouso está Independentemente Associada aos Escores de Classificação de Gordura Visceral em Homens Adultos Sauditas. Arq Bras Cardiol. 2024;121(5):e20220780. doi: 10.36660/abc.20220780.

