

Chronic Chagasic Cardiomyopathy: Influence of Physical Exercise

Gustavo Augusto Ferreira Mota,¹ Mariana Gatto,¹ Vitor Loureiro da Silva,² Paula Felipe Martinez,² Marina Politi Okoshi¹

Departamento de Clínica Médica – Faculdade de Medicina de Botucatu – Universidade Estadual Paulista (UNESP),¹ São Paulo, SP – Brazil

Instituto Integrado de Saúde – Universidade Federal de Mato Grosso do Sul (UFMS),² Campo Grande, MS – Brazil

Short Editorial related to the article: Exercise Training Reduces Inflammation and Fibrosis and Preserves Myocardial Function and Perfusion in a Model of Chronic Chagas Cardiomyopathy

Chronic Chagas cardiomyopathy results from infection with *Trypanosoma cruzi*, transmitted by triatomine insects. The disease, endemic in Latin America, affects millions of individuals worldwide. In recent decades, the increase in migration has elevated the number of infected people in non-endemic regions from Europe and the United States. Chagas disease remains a neglected disease, characterized by difficulty in early diagnosis and poorly understood pathogenesis.¹

T. cruzi infects cardiomyocytes, immune cells, and fibroblasts, activating an inflammatory response that can reduce the parasite load and cause the death of infected cells. The parasite induces the production of chemokines and cytokines that increase the microbicidal capacity of macrophages and stimulate nitric oxide synthases and the production of reactive oxygen species. CD4 and CD8 T lymphocytes also participate in the inflammatory response with the production of IFN- γ and degranulation of cytotoxic T lymphocytes that lysis compromised cells.²

Although the initial inflammatory response is protective, its perpetuation and amplification lead to a progressive increase in myocardial injury and adverse cardiac remodeling. The acute condition is followed by a reparative process with activation of M2-type macrophages and production of anti-inflammatory interleukins and cytokines from the TGF- β family, which induce fibroblast differentiation and extracellular matrix formation.³ Clinically, patients with chronic chagasic heart disease may present arrhythmias, ventricular dysfunction, and heart failure.

The practice of physical exercise is recommended to improve cardiac remodeling and heart failure from different etiologies.^{4,5} Beneficial effects have been described in

chronic chagasic cardiomyopathy, such as improvement in maximal oxygen consumption, left ventricular (LV) ejection fraction, respiratory muscle strength, microvascular function, and quality of life.^{6,7} In experimental studies, physical exercise, performed before or after acute *T. cruzi* infection, modulated inflammatory reaction, improved resistance to *T. cruzi*, and reduced serum concentration of myocardial creatine kinase and myocardial fibrosis.^{8,9} In chronic disease, low-intensity aerobic exercise improved myocardial morphological parameters.¹⁰

In the current issue of ABC Cardiol, the study by Damasceno et al.¹¹ revealed interesting data on the effects of aerobic physical training on myocardial perfusion and LV morphological and functional variables in Syrian hamsters with chronic Chagas cardiomyopathy. Myocardial perfusion defects worsened over time and were associated with reduced LV ejection fraction, increased myocardial inflammation, and Type I collagen expression. A reduction in the gastrocnemius cross-sectional area, characterizing skeletal muscle atrophy, was also observed. Aerobic training attenuated LV systolic dysfunction, perfusion changes, myocardial inflammation, and skeletal muscle atrophy. As pointed out by the authors, only resting perfusion defects were analyzed in the study. Evaluation of perfusion changes under stress situations could bring valuable information.

The effect of physical exercise on Chagas cardiomyopathy is still an open research area. Future studies analyzing the influence of different modalities of exercise and the molecular mechanisms involved in the beneficial effects of exercise on chronic Chagas cardiomyopathy will shed more light on this issue.

Keywords

Cardiomyopathies; Chagas Disease; Inflammation; Functional Status; Exercise; Myocardial Perfusion Imaging

Mailing Address: Gustavo Augusto Ferreira Mota •

Universidade Estadual Paulista Julio de Mesquita Filho – Av. Prof. Mário Rubens Guimarães Montenegro, S/N. Postal Code 18618-687, Botucatu, SP – Brazil

E-mail: gustavo.mota@unesp.br

Manuscript received July 23, 2024, revised manuscript August 14, 2024, accepted August 14, 2024

DOI: <https://doi.org/10.36660/abc.20240507i>

References

1. Padilla AM, Wang W, Akama T, Carter DS, Easom E, Freund Y, et al. Discovery of an Orally Active Benzoxaborole Prodrug Effective in the Treatment of Chagas Disease in Non-human Primates. *Nat Microbiol.* 2022;7(10):1536-46. doi: 10.1038/s41564-022-01211-y.
2. Rigoni TS, Vellozo NS, Guimarães-Pinto K, Cabral-Piccin M, Fabiano-Coelho L, Matos-Silva TC, et al. Axl Receptor Induces Efferocytosis, Dampens M1 Macrophage Responses and Promotes Heart Pathology in Trypanosoma Cruzi Infection. *Commun Biol.* 2022;5(1):1421. doi: 10.1038/s42003-022-04401-w.
3. Maldonado E, Rojas DA, Urbina F, Solari A. The Oxidative Stress and Chronic Inflammatory Process in Chagas Disease: Role of Exosomes and Contributing Genetic Factors. *Oxid Med Cell Longev.* 2021;2021:4993452. doi: 10.1155/2021/4993452.
4. Mota GAF, Souza SLB, Vileigas DF, Silva VL, Sant'Ana PG, Costa LCDS, et al. Myocardial Proteome Changes in Aortic Stenosis Rats Subjected to Long-term Aerobic Exercise. *J Cell Physiol.* 2024;239(4):e31199. doi: 10.1002/jcp.31199.
5. Guizoni DM, Oliveira-Junior SA, Noor SL, Pagan LU, Martinez PF, Lima AR, et al. Effects of Late Exercise on Cardiac Remodeling and Myocardial Calcium Handling Proteins in Rats with Moderate and Large Size Myocardial Infarction. *Int J Cardiol.* 2016;221:406-12. doi: 10.1016/j.ijcard.2016.07.072.
6. Mendes FSNS, Mediano MFF, Souza FCC, Silva PS, Carneiro FM, Holanda MT, et al. Effect of Physical Exercise Training in Patients with Chagas Heart Disease (from the PEACH STUDY). *Am J Cardiol.* 2020;125(9):1413-20. doi: 10.1016/j.amjcard.2020.01.035.
7. Calderon-Ramirez PM, Fernandez-Guzman D, Caira-Chuquineyra B, Mamani-García CS, Medina HM, Diaz-Arocutipa C. Exercise-based Training Programs for Patients with Chronic Chagas Cardiomyopathy: A Systematic Review and Meta-analysis. *Int J Cardiol Heart Vasc.* 2023;48:101256. doi: 10.1016/j.ijcha.2023.101256.
8. Lucchetti BFC, Zanluqui NG, Raquel HA, Lovo-Martins MI, Tatakihara VLH, Belém MO, et al. Moderate Treadmill Exercise Training Improves Cardiovascular and Nitrergic Response and Resistance to Trypanosoma Cruzi Infection in Mice. *Front Physiol.* 2017;8:315. doi: 10.3389/fphys.2017.00315.
9. Pedra-Rezende Y, Barbosa JMC, Bombaça ACS, Dantas-Pereira L, Gibaldi D, Vilar-Pereira G, et al. Physical Exercise Promotes a Reduction in Cardiac Fibrosis in the Chronic Indeterminate Form of Experimental Chagas Disease. *Front Immunol.* 2021;12:712034. doi: 10.3389/fimmu.2021.712034.
10. Preto E, Lima NE, Simardi L, Fonseca FLA, Fragata AA Filho, Maifrino LBM. Effect of Mild Aerobic Training on the Myocardium of Mice with Chronic Chagas Disease. *Biologics.* 2015;9:87-92. doi: 10.2147/BTT.S85283.
11. Damasceno TR, Tanaka DM, Magnani EF, Oliveira RDB, Pereira DAC, Vieira-Alves I, et al. Treinamento Físico Reduz a Inflamação e a Fibrose e Preserva a Função e a Perfusão Miocárdica em um Modelo de Cardiomiopatia Chagásica Crônica. *Arq Bras Cardiol.* 2024; 121(8):e20230707. DOI: <https://doi.org/10.36660/abc.20230707>.

